arXiv:astro-ph/0012014v1 1 Dec 2000

Λ-Inflation and CMBR Anisotropy

Vladimir N. Lukash

Astro Space Center of P.N.Lebedev Physical Institute
Profsoyuznaya 84/32, Moscow, 117810, Russia

Abstract. We argue that a Λ-inflation model can ensure large relative contribution of cosmic gravity waves into the $\Delta T/T$ at COBE scale preserving at the same time a near scale-invariant spectrum of cosmological density perturbations favored by observational data ($n_S \simeq 1$). High efficiency of these models to meet observational tests is discussed.

INTRODUCTION

Observational reconstruction of the power spectrum of cosmological density perturbations is a key problem of the modern cosmology. It provided a dramatic challenge after detecting the primordial CMB anisotropy by DMR COBE as the signal found at 10^0 appeared to be few times higher than the expected value of $\Delta T/T$ in the most simple CDM model considered for many years as a 'standard' cosmological model. During recent years there were many proposals to improve sCDM by adding cosmological constant, hot dark matter, or considering non-scale-invariant primordial spectra of density perturbations.

Presumably, a more natural way to solve the $\Delta T/T$ problem is an introduction of the cosmic gravitational waves contributing into large-scale CMB anisotropy. An inflationary model providing a sufficient abundance of gravity waves should preserve the near-scale-invariant spectrum of density perturbations as indicated by LSS observations.

A simple model of such kind is Λ-inflation, an inflationary model with an effective metastable Λ-term [1]. We briefly recall the physical properties of Λ-inflation and then consider cosmological implications of one of the model with self-interaction.

PHYSICAL PROPERTIES OF Λ-INFLATION

- Λ-inflation is a class of single scalar field models with a positive residual potential energy at the local minimum point, $V_0 > 0$. The hybrid inflation model (at the intermediate evolutionary stage) is a particular case of Λ-inflation: a relation of hybrid to Λ-inflation is similar to the relation of massive field to chaotic inflation. The chaotic inflation is a measure-zero model in the family of Λ-inflation models.

- The scalar perturbation spectrum generated in Λ-inflation has a non-power-law wing-like shape with a broad minimum at $k = k_{cr}$ where the slope is locally scale invariant ($n_S = 1$); it is blue, $n_S > 1$ (red, $n_S < 1$), on short, $k > k_{cr}$ (large, $k < k_{cr}$), scales. The tensor perturbation spectrum remains always red with a maximum deviation (from the Harrison-Zel'dovich one, $n_T = 0$) at the scale near the S-spectrum minimum.
- The cosmic gravity waves generated in Λ-inflation contribute maximally to the Sachs-Wolfe $\Delta T/T$ anisotropy, $(\mathrm{T/S})_{max} \lesssim 10$, at scale where the S-spectrum is slightly red or nearly HZ ($k \lesssim k_{cr}$). T/S remains small ($\ll 1$) in both blue ($k > k_{cr}$) and far red ($k \ll k_{cr}$) S-spectrum regions.
- Three independent arbitrary parameters determine Λ-inflation model; they can be T/S, k_{cr} (the scale where $n_S = 1$), and $\sqrt{V_0}$ (the T-spectrum amplitude at k_{cr} scale). This gives a high capability for fitting various observational tests in the dark matter cosmologies based on Λ-inflation.
- A simple estimate allows to relate straightforwardly the important parameters of observational and physical cosmology:

$$\frac{\mathrm{T}}{\mathrm{S}} \simeq 3r \simeq -6n_T \simeq 12\gamma \simeq \left(\frac{H}{6\times 10^{13}\mathrm{Gev}}\right)^2 . \tag{1}$$

  The ratio between tensor and scalar power spectra, r, and the inflationary γ and Hubble, H, parameters are estimated in scale where the T/S is measured. A large T/S is expected if $(V_0)^{1/4} \sim 10^{16}$ Gev.

Λ-INFLATION WITH SELF-INTERACTION

A general potential of Λ-inflation can be presented as follows:

$$V(\varphi) = V_0 + \sum_{\kappa=2}^{\kappa_{max}} \frac{\lambda_\kappa}{\kappa}\varphi^\kappa, \tag{2}$$

where φ is the inflaton scalar field, $V_0 > 0$ and λ_κ are constants, and $\kappa = 2, 3, 4, ...$

In the case of a massive inflaton ($\kappa = \kappa_{max} = 2$, $\lambda_2 \equiv m^2 > 0$) T/S can be larger than unity only when the S-spectrum slope in the 'blue' asymptote is very steep, $n_S^{blue} > 1.8$. To avoid such a strong spectral bend on short scales ($k > k_{cr}$), we choose another simple version of Λ-inflation – the case with self-interaction (to be called Λλ-inflation): $\kappa = \kappa_{max} = 4$, $\lambda_4 \equiv \lambda > 0$.

The scalar field φ drives an inflationary evolution if $\gamma \equiv -\dot{H}/H^2 < 1$ where $H \simeq \sqrt{V/3}$ (we assume Planckian units $8\pi G = c = \hbar = 1$). This condition holds true for all values of φ if

$$c \equiv \frac{1}{4}\varphi_{cr}^2 = \frac{1}{2}\sqrt{\frac{V_0}{\lambda}} > 1, \tag{3}$$

The gravitational perturbation spectra q_k and h_k generated in S and T modes, respectively, are as follows [2], [1]:

$$q_k = \frac{H}{2\pi\sqrt{2\gamma}} = \frac{\sqrt{2\lambda/3}}{\pi}\left(c^2+x^2\right)^{3/4}, \quad h_k = \frac{H}{\pi\sqrt{2}} = \frac{2c\sqrt{\lambda/3}}{\pi}\left(1+\frac{x}{\sqrt{c^2+x^2}}\right)^{-1/2}, \tag{4}$$

where

$$x = \ln\left[\frac{k}{k_{cr}}\left(1+\left(\frac{x}{c}\right)^2\right)^{1/4}\left(1+\frac{x}{\sqrt{c^2+x^2}}\right)^{2/3}\right] \simeq \ln(k/k_{cr}). \tag{5}$$

CDM COSMOLOGY FROM $\Lambda\lambda$-INFLATION

Let us consider the S-spectrum with CDM transfer function, $T(k)$, normalized both at the large-scale $\Delta T/T|_{10^0}$ (including the contribution from T mode) and the galaxy cluster abundance at $z = 0$, and find a family of the most realistic S-spectra q_k produced in $\Lambda\lambda$-inflation.

In total, we have three parameters entering the q_k spectrum: λ, c and k_{cr}. Constraining them by two observational tests, we are actually left with only one free parameter (say, k_{cr}) which may be restricted elsewhere by other observations.

To demonstrate explicitly how the three parameters are mutually related, we first employ a simple analytic estimates for the σ_8 and $\Delta T/T$ tests to derive the key equation relating c and k_{cr}, and then solve this equation numerically to obtain the range of interesting physical parameters [3].

Instead of taking the σ-integral numerically we can estimate the spectrum amplitude on cluster scale as

$$q_{k_1} \simeq 4.5\times 10^{-7}\frac{h^2\sigma_8}{k_1^2 T(k_1)} . \tag{6}$$

On the other hand, the spectrum amplitude on large scale ($k_2 = k_{COBE} \simeq 10^{-3}h/\mathrm{Mpc}$) can be derived from $\Delta T/T$ due to the Sachs-Wolfe effect:

$$\langle \left(\frac{\Delta T}{T}\right)^2 \rangle_{10^0} = \mathrm{S}+\mathrm{T} \simeq 1.1\times 10^{-10}, \quad \mathrm{S} = 0.04\,\langle q^2\rangle_{\mathrm{COBE}}. \tag{7}$$

The relationship between the power spectrum at COBE scale and the variance of the q potential averaged in 10^0-angular-scale, involves an effective interval of the spectral wavelengths contributing to the latter:

$$\langle q^2\rangle^{1/2}_{COBE} = f q_{k_2}, \quad f^2 \sim \ln\left(\frac{k_2}{k_{hor}}\right). \tag{8}$$

To estimate T/S we will use the approximation formula (1) for $x_2 = x_{COBE}$:

$$\frac{T}{S} \simeq -6n_T = \frac{6}{\sqrt{c^2+x_2^2}}\left(1 - \frac{x_2}{\sqrt{c^2+x_2^2}}\right). \tag{9}$$

Evidently, both normalizations determine essentially the corresponding q_k amplitudes at the locations of cluster (k_1) and COBE (k_2) scales. Accordingly, the parameters k_1 and f can slightly vary while changing the local spectrum slopes at the respective wavelengths. However, when the deviation of q_k slope from HZ is small we can just identify the parameters k_1 and f for CDM models with their sCDM values:

$$k_1 \simeq 0.3h/\mathrm{Mpc}, \quad \mathrm{f} \simeq 1.26.$$

Finally, we now write down the two normalization conditions as follows (for $h = 0.5$ and negligible baryon abundance):

$$q_{k_1} \simeq 3\times 10^{-5}\sigma_8, \quad q_{k_2} \simeq 4\times 10^{-5}\left(1+\frac{\mathrm{T}}{\mathrm{S}}\right)^{-1/2}.$$

Comparing it with the theoretical spectrum q_k we get the key equation for the relationship between c and k_{cr},

$$\left(\frac{q_{k_1}}{q_{k_2}}\right)^2 \simeq D\left(1+\frac{\mathrm{T}}{\mathrm{S}}\right), \tag{10}$$

Eq.(10) has a clear physical meaning: the ratio of the S-spectral power at cluster and COBE scales is proportional to σ_8^2 and inversly proportional to the fraction of the scalar mode contributing to the large-scale temperature anisotropy variance, S/(S+T).

Eq.(10) provides quite a general constraint for the fundamental inflation spectra in a wide set of dark matter models using only two basic measurement (the cluster abundance and large scale $\Delta T/T$). Actually, all information on the dark matter is contained in the D-coefficient which can be calculated using the same equation (10) for a simple particular spectrum preserving the given model parameters:

$$D \simeq \frac{0.6\sigma_8^2}{1-3.1\Omega_b}, \quad \Omega_b < 0.2. \tag{11}$$

The solution of eq.(10) is found in the plane $x-c$ for $D = 0.2, 0.3, 0.4$ numerically. For the entire range $0.1 < D < 0.5$ it can be analytically approximated with a precision less that 10% as follows [3]:

$$\ln^2\left(\frac{k_0}{k_{cr}}\right) \simeq E\,(c_0 - c)\,(c + c_1)\ , \quad 2 < c \le c_0 \tag{12}$$

$$E \simeq 1, \quad \ln k_0 \simeq 49D^2 + 1.3, \quad c_0 \simeq 61D^2 + 6.2, \quad c_1 \simeq 44D^2 + 4.0.$$

The tensor mode contribution is approximated similarly (k_0 and k_{cr} are measured in units h/Mpc):

$$\frac{\mathrm{T}}{\mathrm{S}} \simeq \frac{2.53 - 4.3D}{(\ln k_{cr} + 4.65)^{2/3}} + \frac{1}{3}\ . \tag{13}$$

DISCUSSION

We have presented a Λ-inflation model predicting a near scale-invariant spectrum of density perturbations and large amount of cosmic gravitational waves. This model is consistent with COBE $\Delta T/T$ and cluster abundance data. The perturbation spectra depend on one free scale-parameter, k_{cr}, which can be found in further analysis by fitting other observational data. At the location of k_{cr}, the S-spectrum transients smoothly from the red ($k < k_{cr}$) to the blue ($k > k_{cr}$) parts.

Recent observational data favor a slightly red S-spectra at large scales (cluster power spectrum, BOOMERanG and MAXIMA-1), and require a blue increase of the spectrum at galaxy and subgalactic scales (Ly_α forest). This can be achieved by adjusting k_{cr} to a cluster scale ($\sim 10\,h^{-1}$ Mpc).

ACKNOWLEDGMENTS

The work was partly supported by INTAS grant (97-1192). The author is grateful to the Organizing Committee for the hospitality and financial support.

REFERENCES

1. Lukash V.N., and Mikheeva E.V., *Int.J.Mod.Phys.A* **15**, 3783 (2000).
2. Lukash V.N. In *Cosmology: The Physics of the Universe*, eds. Robson B.A. et al. (World Scientific, Singapore, 1996).
3. Lukash V.N., Mikheeva E.V., M"uller V., Malinovsky A.M., *MNRAS* **317**, 795 (2000).